


**PerCP/Cy5.5 anti-human CD11c**

<b>Catalog # / Size:</b>	2457585 / 25 tests 2457590 / 100 tests
<b>Clone:</b>	S-HCL-3
<b>Isotype:</b>	Mouse IgG2b, $\kappa$
<b>Immunogen:</b>	Spleen cells from patient diagnosed with hairy cell leukemia.
<b>Reactivity:</b>	Human
<b>Preparation:</b>	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
<b>Concentration:</b>	Lot-specific



Human peripheral blood granulocytes were stained with CD11c (clone S-HCL-3) PerCP/Cy5.5 (filled histogram) or mouse IgG2b,  $\kappa$  PerCP/Cy5.5 isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunohistochemistry on frozen tissue sections<sup>1,2,3,4</sup> and immunoprecipitation<sup>1</sup>.

**Application References:**

1. Schwarting R, *et al.* 1985. *Blood* 65:974.
2. Knowles DM, *et al.* 1990. *Am. J. Pathol.* 136:29.
3. Vandenabeele S, *et al.* 2001. *Blood* 97:1733.
4. Shaw JL, *et al.* 2011. *J. Reprod. Immunol.* 89:84.

**Description:** CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin  $\alpha_x$  and CR4. CD11c non-covalently associates with integrin  $\beta_2$  (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing through its interactions with fibrinogen, CD54, and iC3b.

**Antigen References:**

1. Petty HR, Todd RF 3rd. 1996. *Immunol. Today* 17:209.
2. Springer T. 1994. *Cell* 76:301.
3. Ihanus E, *et al.* 2007. *Blood* 109:802-10.