

**PE/Dazzle™ 594 anti-human TIGIT (VSTM3)**

**Catalog # / Size:** 2463575 / 25 tests  
2463580 / 100 tests

**Clone:** A15153G

**Isotype:** Mouse IgG2a, κ


**Immunogen:** Recombinant Human TIGIT.

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



Human peripheral blood leukocytes were stained with CD3 FITC and TIGIT (clone A15153G) PE/Dazzle™ 594 (top) or mouse IgG2a, κ PE/Dazzle™ 594 isotype control (bottom). Data shown was gated on the lymphocyte population.

**Applications:**


**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** This clone can suppress anti-CD3 induced T cell proliferation *in vitro*.



---

**Description:** T cell immunoreceptor with Ig and ITIM domains (TIGIT), also known as VSTM3 or WUCAM, is a 26 kD, type I transmembrane protein and is a member of the PVR (poliovirus receptor) family of immunoglobulin-like domain containing proteins. TIGIT is expressed on activated T cells, follicular T helper, memory, and regulatory T cells as well as on NK cells. TIGIT is a negative regulator of NK and T cell activation. Expression of TIGIT is associated with decreased functionality of CD8 T cells in chronic viral infection and tumors. TIGIT also promotes the differentiation of tolerogenic phenotype in dendritic cells with an increased secretion of IL-10 and a diminished production of IL-12.

**Antigen**  
**References:**

1. Stanietsky N, *et al.* 2009. *Proc. Natl. Acad. Sci.* 106:17858.
2. Yu X, *et al.* 2009. *Nat. Immunol.* 10:48.
3. Johnston R, *et al.* 2014. *Cancer Cell.* 26:923.