

**PE/Cy7 anti-human CD110**

**Catalog # / Size:** 2568550 / 100 tests  
2568545 / 25 tests

**Clone:** S16017E

**Isotype:** Mouse IgG2a, κ


**Immunogen:** Human CD110 transfectants

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



Resting human platelets were stained with PE/Cy7 anti-human CD110 (clone S16017E; filled histogram) or with PE/Cy7 mouse IgG2a

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:**

1. Ninos JM, et al. 2006. J. Transl. Med. 16:4-9.
2. Broudy VC, et al. 1996. Blood. 88:2026-32.
3. Wang X, et al. 2016. Blood. 127:3398-409.

**Description:** CD110 is a type I transmembrane glycoprotein which functions as a receptor for thrombopoietin; binding causes the proliferation and differentiation of megakaryocytes in addition to the production of platelets and protection of stem cells from apoptosis. CD110 is primarily expressed on hematopoietic stem cells, hematopoietic precursor cells, cells of the megakaryocytic lineage, and platelets.

**Antigen References:**

1. Ninos JM, et al. 2006. J. Transl. Med. 16:4-9.
2. Broudy VC, et al. 1996. Blood. 88:2026-32.
3. Wang X, et al. 2016. Blood. 127:3398-409.