

PE/Dazzle™ 594 anti-human CD110

Catalog # / Size: 2569025 / 25 tests
2569030 / 100 tests

Clone: S16017A

Isotype: Mouse IgG2b, κ


Immunogen: Human CD110 transfectants

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Resting human platelets were stained with PE/Dazzle™ 594 anti-human CD110 (clone S16017A; filled histogram) or with PE/Dazzle™ 594 mouse IgG2b, κ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application References:

1. Ninos JM, et al. 2006. J. Transl. Med. 16:4-9.
2. Broudy VC, et al. 1996. Blood. 88:2026-32.
3. Wang X, et al. 2016. Blood. 127:3398-409.

Description: CD110 is a type I transmembrane glycoprotein which functions as a receptor for thrombopoietin; binding causes the proliferation and differentiation of megakaryocytes in addition to the production of platelets and protection of stem cells from apoptosis. CD110 is primarily expressed on hematopoietic stem cells, hematopoietic precursor cells, cells of the megakaryocytic lineage, and platelets.

Antigen

References:

1. Ninos JM, et al. 2006. J. Transl. Med. 16:4-9.
2. Broudy VC, et al. 1996. Blood. 88:2026-32.
3. Wang X, et al. 2016. Blood. 127:3398-409.