## KIRAVIA Blue 520<sup>™</sup> anti-human CD123

Catalog # / Size:	2583550 / 100 tests 2583545 / 25 tests	
Clone:	S18016F	10 <sup>0</sup>
lsotype:	Mouse IgG1, κ	
Immunogen:	Hu CD123 transfectants	
<b>Reactivity:</b>	Human	
Preparation:	The antibody was purified by affinity chromatography and conjugated with KIRAVIA Blue 520™ under optimal conditions.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Human peripheral blood lymphocytes were stained with anti-human CD45RO APC and anti-human CD123 KIRAVIA Blue 520™ (clone S18016F) (right) or mouse IgG1, ĸ KIRAVIA Blue 520™ isotype control (left).
Workshop Number:	IV 103	
Concentration:	Lot-specific	

## **Applications:**

Applications:	Flow Cytometry	
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ L per million cells in 100 $\mu$ L staining volume or 5 $\mu$ L per 100 $\mu$ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.	
	* KIRAVIA Blue 520™ has an excitation maximum of 495 nm, and a maximum emission of 520 nm.	
Application Notes:	S16013D clone can be used for both surface and intracellular detection of TLR9. ICFC compatible with both the intracellular flow cytometric staining and True-Nuclear™ transcription buffer set. Does not work for WB (tested on Daudi cell line).	
Application References:	1. Schreeder DM, <i>et al.</i> 2008. <i>Eur. J. Immunol</i> . 38:3159. (FC)	
Description:	CD123 is the 70 kD transmembrane $\alpha$ chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common $\beta$ chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the $\beta$ chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.	
Antigen References:	1. Miyajima A, <i>et al.</i> 1993. <i>Blood</i> 82:1960.	

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com