

**Alexa Fluor® 700 anti-human CD38**

**Catalog # / Size:** 2586030 / 100 tests  
2586025 / 25 tests

**Clone:** S17015F

**Isotype:** Mouse IgG2a, κ

**Immunogen:** Human CD38 transfectants


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 700 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

**Workshop Number:** IV A053

**Concentration:** Lot-specific



Human peripheral blood lymphocytes were stained with anti-human CD38 (clone S17015F) Alexa Fluor® 700 (filled histogram) or mouse IgG2a, κ Alexa Fluor® 700 isotype control (open histogram).

**Applications:**


**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

**Application Notes:** S17015F is able to cross-block binding of clones HIT2 and HB-7 also raised against human CD38, but not S17015A based on in-house testing.

**Application References:** 1. Li H and Pauza CD. 2015. *Eur. J. Immunol.* 45:298. (IHC)



Human peripheral blood lymphocytes were stained with CD4 APC and CD25 (clone M-A251) KIRAVIA Blue 520™ (left) or mouse IgG1, κ KIRAVIA Blue 520™ isotype control (right).

**Description:** CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

**Antigen** 1. Ferrero E, *et al.* 1999. *J Leuko Biol.* 65:151.  
**References:** 2. Lund F, *et al.* 1995. *Immunol. Today* 16:469.