

PE/Dazzle™ 594 anti-human CD38

Catalog # / Size: 2586095 / 25 tests
2586100 / 100 tests

Clone: S17015F

Isotype: Mouse IgG2a, κ


Immunogen: Human CD38 transfectants

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with anti-human CD38 PE/Dazzle™ 594 (clone S17015F) (filled histogram) or mouse IgG2a, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: S17015F is able to cross-block binding of clones HIT2 and HB-7 also raised against human CD38, but not S17015A based on in-house testing.

Application References: 1. Bühring HJ, *et al.* 1995. *Blood* 86:1916.

Description: CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

Antigen References: 1. Ferrero E, *et al.* 1999. *J Leuko Biol.* 65:151.
2. Lund F, *et al.* 1995. *Immunol. Today* 16:469.