## SONY

## **Product Data Sheet**

## Purified Mouse IgG1, κ Isotype Ctrl

**Catalog # / Size:**  $2600505 / 50 \mu g$ 

2600510 / 500 µg

Clone: MOPC-21

**Isotype:** Mouse IgG1, κ

**Preparation:** The immunoglobulin was purified by

affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.5

## **Applications:**

**Applications:** Other

Recommended Usage:

**d** Each lot of this antibody is quality control tested by immunofluorescent staining

with flow cytometric analysis as negative control. Use at concentrations

comparable to those of the specific antibody of interest.

Application

Notes:

The MOPC-21 immunoglobulin is useful as an isotype-matched control (for the

relevant formats) for Western blotting, immunoprecipitation,

immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF  $^{\text{TM}}$  purified antibody (Endotoxin <0.1 EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for functional assays (Cat. No. 400124) as negative control. For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF  $^{\text{TM}}$  purified antibody (Cat. No. 400166) with a lower endotoxin limit than standard LEAF  $^{\text{TM}}$  purified antibodies (Endotoxin <0.01

EU/microg).

Application References:

- 1. Carlsten M, et al. 2007. Cancer Res. 67:1317. PubMed
- 2. Smed-Sörensen A, et al. 2008. Blood 111:5037. PubMed (FA)
- 3. Bunesmann MM, et al. 2011. Am. J. Respir. Cell. Mol. Biol. Epub. PubMed
- Matsuyama T, et al. 2005. Infect. Immun. 73:1044. (IF)
  Correia DV, et al. 2011. Blood 118:992. (FC) PubMed
- 6. Lian IA, *et al.* 2011. *Placenta*. 32:823. <u>PubMed</u>
- 7. Bufe B, et al. 20015. / Biol Chem. 290:7369. PubMed

**Description:** The MOPC-21 immunoglobulin has unknown specificity. The isotype of this

antibody is mouse IgG1, κ. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human

tissues.