

Brilliant Violet 750™ Mouse IgG1, κ Isotype Ctrl**Catalog # /** 2600530 / 100 tests**Size:** 2600525 / 25 tests**Clone:** MOPC-21**Isotype:** Mouse IgG1, κ**Reactivity:** Mouse**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 750™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 750™ and unconjugated antibody.**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).**Concentration:** Lot-specific**Applications:****Applications:** Flow Cytometry, Intracellular Flow Cytometry**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 750™ excites at 405 nm and emits at 750 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 750™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The MOPC-21 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis.**Application References:**

Description: The MOPC-21 immunoglobulin has unknown specificity. The isotype of this antibody is mouse IgG1, κ . This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human tissues.