

**FITC Mouse IgG1, κ Isotype Ctrl**

**Catalog # / Size:** 2600535 / 50 µg  
2600540 / 200 µg

**Clone:** MOPC-21

**Isotype:** Mouse IgG1, κ

**Preparation:** The immunoglobulin was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis as negative control. Use at concentrations comparable to those of the specific antibody of interest.

**Application Notes:** The MOPC-21 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 400124) as negative control. For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 400166) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

**Application References:**

1. Carlsten M, *et al.* 2007. *Cancer Res.* 67:1317. [PubMed](#)
2. Smed-Sørensen A, *et al.* 2008. *Blood* 111:5037. [PubMed](#) (FA)
3. Bunesmann MM, *et al.* 2011. *Am. J. Respir. Cell. Mol. Biol.* Epub. [PubMed](#)
4. Matsuyama T, *et al.* 2005. *Infect. Immun.* 73:1044. (IF)
5. Correia DV, *et al.* 2011. *Blood* 118:992. (FC) [PubMed](#)
6. Lian IA, *et al.* 2011. *Placenta.* 32:823. [PubMed](#)
7. Bufe B, *et al.* 20015. *J Biol Chem.* 290:7369. [PubMed](#)

---

**Description:** The MOPC-21 immunoglobulin has unknown specificity. The isotype of this antibody is mouse IgG1, κ. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human tissues.