Product Data Sheet

APC Mouse IgG1, κ Isotype Ctrl

Catalog # / Size: 2600595 / 25 µg

2600600 / 100 µg

MOPC-21 Clone:

Isotype: Mouse IgG1, κ

The immunoglobulin was purified by **Preparation:**

> affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated

antibody.

Formulation: Phosphate-buffered solution, pH 7.2.

containing 0.09% sodium azide.

Concentration: 0.2

Applications:

Applications: Flow Cytometry

Recommended

Each lot of this antibody is quality control tested by immunofluorescent staining **Usage:**

with flow cytometric analysis as negative control. Use at concentrations

comparable to those of the specific antibody of interest.

Application

Notes:

The MOPC-21 immunoglobulin is useful as an isotype-matched control (for the

relevant formats) for Western blotting, immunoprecipitation,

immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF $^{\text{\tiny TM}}$ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 400124) as negative control. For in vivo studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 400166) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01

EU/microg).

Application References: 1. Carlsten M, et al. 2007. Cancer Res. 67:1317. PubMed

2. Smed-Sörensen A, et al. 2008. Blood 111:5037. PubMed (FA)

3. Bunesmann MM, et al. 2011. Am. J. Respir. Cell. Mol. Biol. Epub. PubMed

4. Matsuyama T, et al. 2005. Infect. Immun. 73:1044. (IF)

5. Correia DV, et al. 2011. Blood 118:992. (FC) PubMed

6. Lian IA, et al. 2011. Placenta. 32:823. PubMed

7. Bufe B, et al. 20015. J Biol Chem. 290:7369. PubMed

Description: The MOPC-21 immunoglobulin has unknown specificity. The isotype of this

> antibody is mouse IgG1, κ. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human

tissues.