Product Data Sheet

Brilliant Violet 785™ Mouse IgG1, κ Isotype Ctrl

Catalog # / Size: 2600850 / 100 tests

2600845 / 25 tests

Clone: MOPC-21

Isotype: Mouse IgG1, κ

Preparation: The immunoglobulin was purified by

affinity chromatography and conjugated with Brilliant Violet $785^{\,\text{\tiny M}}$ under optimal conditions. The solution is free of unconjugated Brilliant Violet $785^{\,\text{\tiny M}}$ and

unconjugated immunoglobulin.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and BSA

(origin USA).

Concentration: Lot-specific

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis as negative control. Use at concentrations

comparable to those of the specific antibody of interest.

Brilliant Violet 785™ excites at 405 nm and emits at 785 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 785™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes:

The MOPC-21 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF $^{\text{TM}}$ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 400124) as negative control. For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF $^{\text{TM}}$ purified antibody (Cat. No. 400166) with a lower endotoxin limit than standard LEAF $^{\text{TM}}$ purified antibodies (Endotoxin <0.01 EU/microg).

_0,....

Application References:

- 1. Carlsten M, et al. 2007. Cancer Res. 67:1317. PubMed
- 2. Smed-Sörensen A, et al. 2008. Blood 111:5037. PubMed (FA)
- 3. Bunesmann MM, et al. 2011. Am. J. Respir. Cell. Mol. Biol. Epub. PubMed
- 4. Matsuyama T, et al. 2005. Infect. Immun. 73:1044. (IF)
- 5. Correia DV, et al. 2011. Blood 118:992. (FC) PubMed
- 6. Lian IA, et al. 2011. Placenta. 32:823. PubMed

Description:

The MOPC-21 immunoglobulin has unknown specificity. The isotype of this antibody is mouse IgG1, κ . This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human tissues.