Product Data Sheet

PE Mouse IgG2a, κ Isotype Ctrl (FC)

Catalog # / Size: 2601070 / 100 tests

2601065 / 25 tests

Clone: MOPC-173

Isotype: Mouse IgG2a, κ

Preparation: The immunoglobulin was purified by

affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated

immunoglobulin.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this mouse IgG2a, κ isotype control antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, use the isotype control at the same concentration as your primary antibody. Use our <u>Concentration Lookup</u> tool to find the exact concentrations of your lots of product.

Application Notes:

The MOPC-173 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation,

immunohistochemistry, functional assay, and immunofluorescence microscopy. The LEAF[™] purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 400224) as negative control. For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF[™] purified antibody (Cat. No. 400264) with a lower endotoxin limit than standard LEAF[™] purified antibodies (Endotoxin <0.01 EU/microg).

Application References:

- 1. Luckashenak NA, et al. 2006. J. Immunol. 177:5177.
- eferences: 2. Burman AC, et al. 2007. Blood 110:1064.
 - 3. Goo SY, et al. 2007. J. Biol. Chem. doi:10.1074/jbc.M701876200.
 - 4. Podolin PL, et al. 2008. J. Immunol. 180:7989. PubMed
 - 5. Ohno Y, et al. 2013. J Biochem. 154:355. PubMed

Description: The MOPC-173 immunoglobulin has unknown specificity. The isotype of this

antibody is mouse IgG2a, κ. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human

tissues.