

Brilliant Violet 605™ Mouse IgG2b, κ Isotype Ctrl

| | |
|--------------------------|--|
| Catalog # / Size: | 2601750 / 100 tests 2601745 / 25 tests |
| Clone: | MPC-11 |
| Isotype: | Mouse IgG2b, κ |
| Preparation: | The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 605™ and unconjugated antibody. |
| Formulation: | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA). |
| Concentration: | Lot-specific |

Applications:

| | |
|---------------------------|---|
| Applications: | Flow Cytometry |
| Recommended Usage: | Each lot of this mouse IgG2b, κ isotype control antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, use the isotype control at the same concentration as your primary antibody. Use our Concentration Lookup tool to find the exact concentrations of your lots of product. |

Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

| | |
|---------------------------|--|
| Application Notes: | The MPC-11 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 400324) as negative control. For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 400348) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg). |
|---------------------------|--|

| | |
|--------------------------------|---|
| Application References: | 1. Smed-Sørensen A, <i>et al.</i> 2008. <i>Blood</i> 111:5037. (FA) PubMed 2. Podolin PL, <i>et al.</i> 2008. <i>J. Immunol.</i> 180:7989. (FC) PubMed |
|--------------------------------|---|

Description: The MPC-11 immunoglobulin has unknown specificity. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human tissues.