## **Product Data Sheet**

## Brilliant Violet 711™ Mouse IgG2b, κ Isotype Ctrl

Catalog # / Size: 2601770 / 100 tests

2601765 / 25 tests

Clone: MPC-11

**Isotype:** Mouse IgG2b, κ

Preparation: The antibody was purified by affinity

chromatography and conjugated with Brilliant Violet 711<sup>™</sup> under optimal conditions. The solution is free of unconjugated Brilliant Violet 711<sup>™</sup> and

unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and BSA

(origin USA).

Concentration: Lot-specific

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** by immur staining, or

Each lot of this mouse IgG2b,  $\kappa$  isotype control antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, use the isotype control at the same concentration as your primary antibody. The bandpass filter 710/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 711 $^{\text{TM}}$  is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes:

The MPC-11 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 400324) as negative control. For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 400348) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01

EU/microg).

**Application** 1. Smed-Söre 2. Podolin PL,

1. Smed-Sörensen A, *et al.* 2008. *Blood* 111:5037. (FA) <u>PubMed</u> 2. Podolin PL, *et al.* 2008. *J. Immunol.* 180:7989. (FC) <u>PubMed</u>

**Description:** The MPC-11 immunoglobulin has unknown specificity. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and

fixed mouse, rat and human tissues.