Product Data Sheet

PerCP/Cy5.5 Rat IgG2a, κ Isotype Ctrl

Catalog # / Size: $2602655 / 25 \mu g$

2602660 / 100 µg

Clone: RTK2758

Isotype: Rat IgG2a, κ

Immunogen: Trinitrophenol + KLH

Reactivity: Other

Preparation: The immunoglobulin was purified by

affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated immunoglobulin.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2

Applications:

Applications: Flow Cytometry

Recommended

Usage: with flow

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis as negative control. Use at concentrations

comparable to those of the specific antibody of interest.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of

690 nm.

Application Notes:

The RTK2758 immunoglobulin is useful as an isotype-matched control (for the

relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, immunofluorescence microscopy,

immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF $^{\text{\tiny IM}}$ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 400516) as negative control. For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF $^{\text{\tiny IM}}$ purified antibody (Cat. No. 400544) with a lower endotoxin limit than standard LEAF $^{\text{\tiny IM}}$ purified antibodies (Endotoxin <0.01

EU/microg).

Application

1. Nishimoto H, et al. 2005. Blood 106:4241.

References:

2. Seach N, et al. 2008. J. Immunol. 180:5384. PubMed

3. Jiang P, et al. 1999. J. Biol. Chem. 274:559. (FA)

Description: The RTK2758 immunoglobulin reacts with KLH. The isotype of this antibody is rat

IgG2a, K. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human tissues.