## **Product Data Sheet**

## Spark Blue<sup>™</sup> 550 Rat IgG2b, κ Isotype Ctrl

**Catalog #** /  $2603405 / 25 \mu g$ 

**Size:**  $2603410 / 100 \mu g$ 

Clone: RTK4530

**Isotype:** Rat IgG2b, κ

**Immunogen:** Trinitrophenol + KLH

**Preparation:** The antibody was purified by affinity

chromatography and conjugated with Spark Blue™ 550 under optimal

conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide

Concentration: 0.5 mg/mL

## **Applications:**

**Applications:** Flow Cytometry, Intracellular Staining for Flow Cytometry

Recommended Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis as negative control. Use at concentrations comparable to those of the specific antibody of interest. Use

our Concentration Lookup tool to find the exact concentrations of your lots of product.

\* Spark Blue™ 550 has a maximum excitation of 516 nm and a maximum emission of 540 nm.

Application Notes:

The RTK4530 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, and immunofluorescence microscopy. The Ultra-LEAF  $^{\text{TM}}$  purified antibody (Endotoxin < 0.01 EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for functional assays (Cat. No. 400643, 400644, 400671, 400672, 400675, and 400676) as negative

control.

Application References:

- 1. Cervantes-Barragan L, et al. 2007. Blood 109:1131.
- 2. Zeiser R, et al. 2007. Blood 109:2225.
- 3. Sasaki K, et al. 2008. J. Immunol. 181:104. PubMed
- 4. Duan J, et al. 2008. P. Natl. Acad. Sci. USA 105:5183. PubMed
- 5. Yi H, et al. 2009. Blood 113:5819. PubMed
- 6. Schafeer JS, et al. 2010. J. Leukocyte Biol. 87:301. PubMed
- 7. Lei GS, et al. 2015. Infect Immun. 83:572. PubMed
- 8. Richards J, et al. 2015. Mol Cell Cardiol. 79:21. PubMed

**Description:** The isotype of RTK4530 immunoglobulin is rat IgG2b, κ. This antibody was

chosen as an isotype control after screening on a variety of resting,

activated, live, and fixed mouse, rat and human tissues.