Product Data Sheet

Biotin Armenian Hamster IgG Isotype Ctrl

Catalog # / Size: 2604520 / 200 µg

2604515 / 50 μg

Clone: **HTK888**

Isotype: Hamster IgG

Trinitrophenol + KLH Immunogen:

Preparation: The immunoglobulin was purified by

affinity chromatography, and

conjugated with biotin under optimal conditions. The solution is free of

unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.5

Applications:

Applications: Flow Cytometry

Recommended

Each lot of this antibody is quality control tested by immunofluorescent staining **Usage:**

with flow cytometric analysis as negative control. Use at concentrations

comparable to those of the specific antibody of interest.

Application Notes:

The HTK888 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, functional assay, immunofluorescence microscopy, immunocytochemistry and immunofluorescent staining (surface or intracellular) for flow cytometric analysis. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 400916) as negative control. For in vivo studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 400940) with a lower endotoxin limit than standard LEAF™ purified antibodies

(Endotoxin < 0.01 EU/microg).

Application References: 1. Lesley R, et al. 2006. P. Natl. Acad. Sci. USA 103:10717.

2. Yu R, et al. 2006. Obesity 14:1353.

3. Yang JH, et al. 2005. Rheumatology (Oxford). 44:1245. PubMed 3. Mina-Osorio P, et al. 2008. J. Leukocyte Biol. 84:448. PubMed

4. Shen H, et al. 2009. J. Am Soc Nephrol. 20:1032. PubMed

5. Peraino JS, et al. 2013. Hum Immunol. 18:88. PubMed.

Description: This antibody was chosen as an isotype control after screening on a variety of

resting, activated, live, and fixed mouse, rat and human tissues.