## PerCP/Cy5.5 Goat anti-mouse IgG (minimal x-reactivity)

Catalog # / Size: 2626570 / 100 µg

> Clone: Polv4053 Isotype: Goat IgG Reactivity: Mouse

**Preparation:** The antibody was purified by affinity

> chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated


antibody.

Phosphate-buffered solution, pH 7.2, Formulation:

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

**Concentration:** 0.2



Human peripheral blood lymphocytes stained with purified mouse anti-human CD3 (UCHT1), followed by anti-mouse IaG (Poly4053) PerCP/Cy5.5

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of

690 nm.

**Application** Notes: This polyclonal goat anti-mouse IgG antibody is useful for capture or detection of

mouse IgG in ELISA.

**Application** References:

1. Scotta, C, et al. 2008. J. Immunol. 181:1025. PubMed

**Description:** This polyclonal goat anti-mouse IgG antibody reacts with the heavy chains of

mouse IgG and with the light ( $\kappa$  and  $\lambda$ ) chains common to most mouse immunoglobulins. No cross-reactivity has been detected against non-

immunoglobulin serum proteins. This antibody has been solid-phase absorbed to ensure minimal cross-reaction with rat, human, bovine, horse, and rabbit

immunoglobulins, but it may has minimal cross-reaction with other subclasses of

mouse immunoglobulins.