

---

**DyLight™ 649 Goat anti-hamster (Armenian) IgG**

**Catalog # / Size:** 2627525 / 100 µg  
**Clone:** Poly4055  
**Isotype:** Goat IgG  
**Reactivity:** Other  
**Preparation:** The antibody was purified by affinity chromatography, and conjugated with DyLight™ 649 under optimal conditions.  
**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).  
**Concentration:** 0.5

**Applications:**

**Applications:** Immunofluorescence

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25$  microg per  $10^6$  cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* DyLight™ 649 has a maximum absorption of 655nm and a maximum emission of 670nm (similar to Alexa Fluor® 647 and APC).

**Application Notes:** This conjugated goat anti-hamster IgG antibody is useful for immunofluorescent staining with flow cytometric analysis or microscopy.

**Application References:** 1. Porcherie A, *et al.* 2011. *J. Exp Med.* 11:2225. [PubMed](#)  
2. Kim Yu, *et al.* 2015. *PLoS One.* 10:120294. [PubMed](#)

---

**Description:** This polyclonal antibody reacts with the heavy chains of IgG and the light chains common to most immunoglobulins from Armenian hamster. No cross-reactivity has been detected against non-immunoglobulin serum proteins. This antibody has been solid-phase absorbed to ensure minimal cross-reaction with mouse, rat, human, bovine, and rabbit immunoglobulins, but may cross-react with Armenian hamster IgM and Syrian hamster immunoglobulins.