
Product Data Sheet

PerCP/Cy5.5 anti-mouse IgG1

| Catalog # / Size: | 2633055 / 25 μg 2633060 / 100 μg |
|--------------------|---|
| Clone: | RMG1-1 |
| Isotype: | Rat IgG |
| Immunogen: | Mouse Ig cocktail |
| Reactivity: | Mouse |
| Preparation: | The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody. |
| Formulation: | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. |
| Concentration: | 0.2 |



Human peripheral blood lymphocytes were stained with purified anti-human CD3 (clone UCHT1, filled histogram) or mouse lgG1, κ isotype control (open histogram), followed by anti-mouse lgG1 (clone RMG1-1) PerCP/Cy5.5.

Applications:

| Applications: | Flow Cytometry |
|----------------------------|---|
| Recommended Usage: | Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application. |
| | * PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm. |
| Application References: | 1. Rucci F, <i>et al.</i> 2010. <i>P. Natl. Acad. Sci. USA</i> 107:3024. (FC) <u>PubMed</u> 2. Zhang JG, <i>et al.</i> 2011. <i>Int. Immunopharmacol.</i> 11:1685. <u>PubMed</u> |
| Description: | The RMG1-1 monoclonal antibody reacts with immunoglobulin G1 (IgG1) in all tested mouse haplotype (Igh-a and b). It does not react with other isotypes. The RMG1-1 monoclonal antibody may be used as a primary or secondary reagent for ELISA or immunofluorescent analysis. |