

APC/Fire™ 750 anti-mouse IgG1

Catalog # / Size: 2633115 / 25 µg
2633120 / 100 µg

Clone: RMG1-1

Isotype: Rat IgG


Immunogen: Mouse Ig cocktail

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml



Human peripheral blood lymphocytes were stained with purified anti-human CD3 (clone UCHT1 isotype mouse IgG1, κ) followed by APC/Fire™ 750 anti-mouse IgG1 antibody (filled histogram) or APC/Fire™ 750 anti-mouse IgG1 antibody alone (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested. For flow cytometric staining, the suggested use of this reagent is ≤0.125 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application References: 1. Rucci F, et al. 2010. *P. Natl. Acad. Sci. USA* 107:3024. (FC) [PubMed](#)
2. Zhang JG, et al. 2011. *Int. Immunopharmacol.* 11:1685. [PubMed](#)

Description: The RMG1-1 monoclonal antibody reacts with immunoglobulin G1 (IgG1) in all tested mouse haplotype (Igh-a and b). It does not react with other isotypes. The RMG1-1 monoclonal antibody may be used as a primary or secondary reagent for ELISA or immunofluorescent analysis.