

Biotin anti-mouse IgMa

Catalog # / Size: 2643015 / 50 µg
2643020 / 500 µg

Clone: MA-69


Isotype: Mouse IgG1, κ

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



BALB/c splenocytes stained with MA-69 Biotin, followed by Sav-PE

Applications:

Applications: Other

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10^6 cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: ELISA as capture antibody or detection antibody, stimulation of B cell proliferation, and immunohistochemistry of acetone-fixed frozen sections.

Application References: 1. Sieckmann D, *et al.* 1991. *Hybridoma* 10:121.
2. Stall AM. 1996. Weir's Handbook of Experimental Immunology pp. 27.

Description: This antibody was raised by immunization with IgM^a mAb TC31. It specifically reacts with mouse IgM of *Igh-C^a* and the related haplotypes (c, f, g, h, and j). It has been reported that beads coated with MA-69 antibody can stimulate B cell proliferation.