


Biotin anti-human IgG Fc

Catalog # / Size: 2653590 / 25 µg
Clone: M1310G05
Isotype: Rat IgG2a, κ
Immunogen: Human Siglec-E-IgG Fc fusion protein.
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5




Human peripheral blood lymphocytes were stained with CD19 APC and IgG (clone M1310G05) Biotin (top) or rat IgG2a, κ Biotin isotype control (bottom).

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone M1310G05 recognizes IgG in the membrane of memory B cells, has a stronger affinity for IgG1 and IgG3 than for IgG2 and IgG4, and does not cross react with IgD, IgE, or IgM.

Application References: 1. Paul, WE. (2003). *Fundamental Immunology*. Philadelphia, PA: Lippincott, Williams, & Wilkins.



Description: IgG Fc is a homodimer that is composed of the constant region of the two heavy chains that form the IgG molecule. The Fc fragment mediates opsonization, antibody dependent cellular cytotoxicity (ADCC), and complement activation through binding to Fc receptors such as CD16, CD32, CD64, and the complement factor C1.

Antigen References: 1. Paul, WE. (2003). *Fundamental Immunology*. Philadelphia, PA: Lippincott, Williams, & Wilkins.