


APC/Fire™ 750 anti-human IgG Fc**Catalog # /** 2653620 / 100 tests**Size:** 2653615 / 25 tests**Clone:** M1310G05**Isotype:** Rat IgG2a, κ**Immunogen:** Human Siglec-E-IgG Fc fusion protein.**Reactivity:** Human**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).**Concentration:** Lot-specific

Human peripheral blood lymphocytes were stained with CD19 BV421 and IgG Fc (clone M1310G05) APC/Fire™ 750 (left) or rat IgG2a, κ APC/Fire™ 750 isotype control (right).

Applications:**Applications:** Flow Cytometry**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: Clone M1310G05 recognizes IgG in the membrane of memory B cells, has a stronger affinity for IgG1 and IgG3 than for IgG2 and IgG4, and does not cross react with IgD, IgE, or IgM.**Description:** IgG Fc is a homodimer that is composed of the constant region of the two heavy chains that form the IgG molecule. The Fc fragment mediates opsonization, antibody dependent cellular cytotoxicity (ADCC), and complement activation through binding to Fc receptors such as CD16, CD32, CD64, and the complement factor C1.**Antigen References:** 1. Paul, WE. (2003). *Fundamental Immunology*. Philadelphia, PA: Lippincott, Williams, & Wilkins.