


PerCP/Cy5.5 anti-human IL-2

Catalog # / Size:	3101605 / 25 tests 3101610 / 100 tests
Clone:	MQ1-17H12
Isotype:	Rat IgG2a, κ
Immunogen:	<i>E. coli</i> -expressed recombinant human IL-2
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Concentration:	Lot-specific



PMA + ionomycin-stimulated (6 hours) human peripheral blood lymphocytes intracellular stained with MQ1-17H12 PerCP/Cy5.5 and CD3 (UCHT1) PE

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes:	<p>ELISA or ELISPOT Capture^{2,3}: The purified MQ1-17H12 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated Poly5176 antibody (Cat. No. 517605) as the detecting antibody. The LEAF[™] purified antibody is suggested for ELISPOT capture. For ELISPOT capture applications, a concentration range of 4-8 microg/ml is recommended.</p> <p>Additional reported applications (for the relevant formats) include: immunoprecipitation², immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections^{1,4-6,8}, neutralization¹³, and immunocytochemistry.</p>
---------------------------	---

Note: For testing human IL-2 in serum or plasma, BioLegend's LEGEND MAX[™] Kits (Cat. No. 431807 & 431808) are specially developed and recommended.

Application References:	<ol style="list-style-type: none"> Andersson J, <i>et al.</i> 1994. <i>Immunology</i> 83:16. (IHC) Abrams J, <i>et al.</i> 1992. <i>Immunol. Rev.</i> 127:5. (IP) Abrams JS. 1995. <i>Curr. Prot. Immunol.</i> Unit 6.20. Fernandez V, <i>et al.</i> 1994. <i>Eur. J. Immunol.</i> 24:1808. (IHC) Skansen-Saphir U, <i>et al.</i> 1994. <i>Eur. J. Immunol.</i> 24:916. (IHC) Andersson U, <i>et al.</i> <i>Detection and Quantification of Gene Expression</i>. New York:Springer-Verlag. (IHC)
--------------------------------	--

7. Prussin C, *et al.* 1995. *J. Immunol. Methods*. 188:117.
 8. Raqib R, *et al.* 2002. *Infect. Immun.* 70:3199. (IHC)
 9. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
 10. Colleton BA, *et al.* 2009. *J Virol.* 83:6288. [PubMed](#)
 11. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 12. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
 13. Steindor M, *et al.* 2015. *PLoS One.* 10:19737. [PubMed](#)
-

Description: IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

Antigen
References:

1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.
2. Taniguchi T, *et al.* 1993. *Cell* 73:5.
3. Nistico G. 1993. *Prog. Neurobiol.* 40:463.
4. Waldmann T, *et al.*