

Brilliant Violet 605™ anti-human IL-4

Catalog # / Size: 3104135 / 25 tests
3104140 / 100 tests

Clone: MP4-25D2

Isotype: Rat IgG1, κ


Immunogen: CHO-expressed, recombinant human IL-4

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 605™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific




PMA⁺ ionomycin-stimulated human peripheral blood lymphocytes (6 hours, in the presence of monensin) were surface stained with CD3 APC, fixed, permeabilized and then stained with IL-4 (clone MP4-25D2) Brilliant Violet 605™ (top) or rat IgG (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use

the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes:

ELISA Detection^{1,3} or ELISPOT

Detection^{4,5}: The biotinylated MP4-25D2 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified 8D4-8 antibody (Cat. No. 500702/500707) as the capture antibody.

Flow Cytometry^{6,9}: The fluorochrome-labeled MP4-25D2 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-4-producing cells within mixed cell populations.

Neutralization¹⁻³: The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for neutralization of human IL-4 bioactivity (Cat. No. 500815). The MP4-25D2 antibody can neutralize the bioactivity of natural or recombinant IL-4.

Application References:

1. Chretien I, *et al.* 1989. *J. Immunol. Methods* 117:67. (ELISA Detection, Neut)
2. Ramanathan L, *et al.* 1993. *Biochem.* 32:3549. (Neut)
3. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (ELISA Detection, Neut)
4. Mahanty S, *et al.* 1992. *J. Immunol.* 148:3567. (ELISPOT Detection)
5. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.19. (ELISPOT Detection)
6. Prussin C, *et al.* 1995. *J. Immunol. Methods* 188:117. (ICFC)
7. Raqib R, *et al.* 1995. *Infect. Immun.* 63:289.
8. Andersson J, *et al.* 1994. *Immunology* 83:16.
9. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
10. Kubota M, *et al.* 1997. *J. Immunol.* 158:5321.
11. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
12. Kroneke MA, *et al.* 2012. *J. Immunol.* 188:3734. [PubMed](#)

Description:

IL-4 is a pleiotropic cytokine that is produced by activated T cells, mast cells, and basophils. IL-4 elicits many different biological responses but has two dominant functions. The first is regulating differentiation of naïve CD4⁺ T cell to the Th2 type. Th2 cells produce IL-4, IL-5, IL-10, and IL-13, which tend to favor a humoral immune response while suppressing a cell-mediated immune response controlled by Th1 cells. The second is regulating IgE and IgG1 production by B cells.

Antigen References:

1. Fitzgerald K, *et al.* Eds. 2001. *The Cytokine FactsBook.* Academic Press San Diego.
2. Boulay J, *et al.* 1992. *Curr. Opin. Immunol.* 4:294.
3. Dullens H, *et al.* 1991. *In vivo* 5:567.
4. Paul