

**Brilliant Violet 421™ anti-mouse IL-2**

**Catalog # / Size:** 3119125 / 125 µl  
3119130 / 50 µg

**Clone:** JES6-5H4

**Isotype:** Rat IgG2b, κ


**Immunogen:** *E. coli*-expressed, recombinant mouse IL-2

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

**Concentration:** NULL




C57BL/6 mouse splenocytes were stimulated with PMA + Ionomycin for 6 hours (in the presence of monensin), stained with CD3 FITC, fixed, permeabilized, and then stained with IL-2 (clone JES6-5H4) Brilliant Violet 421™ (top) or rat IgG2b, κ Brill

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by

U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:**

**ELISA Detection<sup>1-3</sup> or ELISPOT**

**Detection<sup>4-6</sup>:** The biotinylated JES6-5H4 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with the purified JES6-1A12 antibody (Cat. No. 503702/503704) as capture antibody and recombinant mouse IL-2 (Cat. No. 575409) as the standard.

**Flow Cytometry<sup>8-10</sup>:** The fluorochrome-labeled JES6-5H4 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-2 - producing cells within mixed cell populations.

**Neutralization<sup>1,7</sup>:** The LEAF™ purified antibody (Endotoxin in vivo and *in vitro* (Cat. No. 503812) is recommended for neutralization.

**Additional reported applications (for the relevant formats) include:**

immunoprecipitation<sup>1</sup>, immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections<sup>2</sup>, *in vivo* capture<sup>7</sup>, and immunocytochemistry.

**Note:** For testing mouse IL-2 in serum, plasma or supernatant, BioLegend's ELISA MAX™ Sets (Cat. No. 431001 to 431006) are specially developed and recommended.

**Application References:**

1. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5.
2. Sander B, *et al.* 1993. *J. Immunol. Meth.* 166:201.
3. Abrams J. 1995. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.20.
4. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.19.
5. Mo X, *et al.* 1995. *J. Virol.* 69:1288.
6. Karulin A, *et al.* 2000. *J. Immunol.* 164:1862.
7. Finkelman F, *et al.* 2003. *Curr. Prot. Immunol.* John Wiley & Sons New York. Unit 6.28.
8. Ko SY, *et al.* 2005. *J. Immunol.* 175:3309. [PubMed](#)
9. Kang SS and Allen PM. 2005. *J. Immunol.* 174:5382.
10. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366.

---

**Description:**

IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

**Antigen References:**

1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.
2. Taniguchi T, *et al.* 1993. *Cell* 73:5.
3. Nistico G. 1993. *Prog. Neurobiol.* 40:463.
4. Waldmann T, *et al.*