

Alexa Fluor® 700 anti-human IFN-γ

Catalog # / Size: 3132575 / 25 µg
3132580 / 100 µg

Clone: B27

Isotype: Mouse IgG1, κ


Immunogen: *E. coli*-expressed recombinant human IFN-γ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 700 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



PMA+ionomycin-stimulated (5 hours) human PBMCs surface stained with CD3 PE and intracellular stained with B27 Alexa Fluor® 700

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is highly recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: **Flow Cytometry²:** The fluorochrome-labeled B27 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN-γ -producing cells within mixed cell populations. For intracellular cytokine staining protocol, please visit www.biolegend.com and click on the support section.

Neutralization^{1,3,6,7}: The LEAF™ Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for neutralization of human IFN-γ bioactivity (Cat. No. 506512).

- Application References:**
1. Favre C, *et al.* 1989. *Molec. Immunol.* 26:17. (Neut)
 2. Kaur A, *et al.* 2002. *J Virol.* 76:3646.
 3. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (Neut)
 4. Andersson U, *et al.* 1999. *Detection and quantification of gene expression.* New York:Springer-Verlag.
 5. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
 6. Acosta-Rodriguez EV, *et al.* 2007. *Nat. Immunol.* 9:942-9. (Neut)
 7. Gangur V, *et al.* 1998. *FASEB J.* 12:705-13. (Neut)

Description: Interferon-γ is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral

activities, IFN- γ also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN- γ can upregulate MHC class I and II antigen expression by antigen-presenting cells. The B27 antibody reacts with the human interferon- γ . The B27 antibody can neutralize the bioactivity of natural or recombinant IFN- γ .

**Antigen
References:**

1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.
2. De Maeyer E, *et al.* 1992. *Curr. Opin. Immunol.* 4:321.
3. Farrar M, *et al.* 1993. *Annu. Rev. Immunol.* 11:571.