

PerCP/Cy5.5 anti-human IL-9

Catalog # / Size: 3138045 / 25 tests
3138050 / 100 tests

Clone: MH9A4

Isotype: Mouse IgG2b, κ


Immunogen: Baculovirus-expressed, recombinant human IL-9

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific




PMA+ionomycin-stimulated (6 hours, in the presence of monensin) Th2-polarized lymphocytes were stained with CD3 APC, permeabilized, and intracellularly stained with IL-9 (clone MH9A4) PerCP/Cy5.5 (top image), or mouse IgG2b, κ PerCP/Cy5.5 isotype co

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: **ELISA Capture2:** The purified MH9A4 antibody is useful as the capture antibody in a human IL-9 sandwich ELISA assay, when used in conjunction with the biotinylated MH9D1 (Cat. No. 507702) antibody as the detecting antibody.

Flow Cytometry: The fluorochrome-labeled MH9A4 antibody is useful for intercellular immunofluorescent staining and flow cytometric analysis to identify human IL-9-producing cells in mixed cell populations. For intracellular cytokine staining protocol, please visit www.biolegend.com and click on the

support section.

- Application** 1. Jenmalm M, *et al.* 2001. *Clin. Exptl. Aller.* 31:1528.
- References:** 2. Faulkner H, *et al.* 2002. *J. Infec. Diseas.* 185:665.
3. Chen J, *et al.* 2008. *Blood* 111:5163. [PubMed](#)
4. Chang HC, *et al.* 2010. *Nat. Immunol.* 11:527. (ELISA) [PubMed](#)
5. Lozano E, *et al.* 2012. *J Immunol.* 188:3869. [PubMed](#).
-

Description: IL-9 is a potent, T cell-derived, T cell growth factor which can also enhance mast cell activity and IL-3- or IL-4- dependent proliferation of bone marrow-derived mast cells. IL-9 synergizes with erythropoietin to promote erythroid colony formation. IL-9 has also been reported to protect human T cells from apoptosis induced by IL-2 withdrawal. IL-9 is upregulated in human eosinophils by TNF- α and IL1- β . IL-9 has been reported to downregulate the oxidative burst in activated human alveolar macropahges and induce TGF- β production. The MH9A4 antibody reacts with human IL-9. The MH9A4 antibody can neutralize the bioactivity of natural or recombinant IL-9.

- Antigen** 1. Fitzgerald K, *et al.* Eds. 2001. *The Cytokine FactsBook.* Academic Press San Diego.
- References:** 2. Quesniaux V. 1992. *Research Immunology* 143:385.
3. Renauld J, *et al.* 1993. *Adv. Immunol.* 54:79.
4. Yang Y. 199