

Alexa Fluor® 647 anti-human IL-1β

Catalog # / 3141040 / 100 tests
Size: 3141035 / 25 tests

Clone: JK1B-1

Isotype: Mouse IgG1, κ

Immunogen: Recombinant human IL-1 beta


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: HCDM listed

Concentration: Lot-specific



LPS-stimulated (6 hour) human peripheral blood monocytes surface stained with CD14 PE, then intracellular stained with JK1B-1 Alexa Fluor® 647

Applications:


Applications: Intracellular Staining for Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

Application Notes: **Other and ELISPOT Capture:** The purified JK1B-1 antibody is useful as the capture antibody in a sandwich Other assay, when used in conjunction with the biotinylated JK1B-2 antibody as the detecting antibody.

Flow Cytometry: The fluorochrome-labeled JK1B-1 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-1β-producing cells within mixed cell populations.



Application References: 1. Loiarro M, et al. 2013. *J Biol Chem*. 288:30210. [PubMed](#)
 2. Fell LH, et al. 2014. *Nephrol Dial Transplant*. [PubMed](#)

Description: IL-1 refers to two proteins, IL-1 α and IL-1 β which are the products of distinct genes, but which are recognized by the same cell surface receptors. IL-1 β is a potent immuno-modulator which mediates a wide range of immune and inflammatory responses including the activation of B and T cells. The JK1B-1 antibody reacts with human interleukin-1 β (IL-1 β) and baboon IL-1 β . The JK1B-1 antibody does not recognize mouse IL-1 β or human IL-1 α or other cytokines. The JK1B-1 antibody can neutralize the bioactivity of natural or recombinant IL-1 β .

Antigen
References:

1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.
2. Bomford R, *et al.* Eds. 1989. *Interleukin-1 inflammation and disease*. Elsevier New York.
3. Brazel D, *et al.* 1991. *Biotechnol. Ther.* 2:241.
4. Dinarello C. 1996. *Blood* 87:2095.