


FITC anti-Cytochrome c

Catalog # / Size: 3661520 / 100 µg
Clone: 6H2.B4
Isotype: Mouse IgG1, κ
Immunogen: Rat cyt c-OVA
Reactivity: Human, Mouse, Rat
Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.
Formulation: This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



Balb/c splenocytes intracellularly stained with 6H2.B4 FITC

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 microg per 10⁶ cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes: Additional reported applications (for the relevant formats) include: intracellular flow cytometry⁵, immunofluorescence microscopy^{3,5}, immunoprecipitation⁴, and immunocytochemistry⁵.
Application References: 1. Goshorn SC, *et al.* 1991. *J. Biol. Chem.* 266:2134.
 2. Jemmerson R, *et al.* 1991. *Eur. J. Immunol.* 21:143.
 3. Chandra D, *et al.* 2002. *J. Biol. Chem.* 277:50842. (IF)
 4. Semenkova L, *et al.* 2003. *Eur. J. Biochem.* 270:4388. (IP)
 5. Shih S-F, *et al.* 2001. *J. Biol. Chem.* 276:21870. (ICFC ICC IF)
 6. Ma Y, *et al.* 2013. *Brain Res.* 1351:222. [PubMed](#)

Description: Cytochrome c is a 15 kD protein found in the mitochondrial intermembrane space with a heme-binding domain. Cytochrome c is a component of the electron transport chain; the heme group transfers electrons from cytochrome b-c1 complex to cytochrome oxidase complex. Cytochrome c initiates apoptosis by release to cytoplasm and binding Apaf-1 which activates procaspase 9. Cytochrome c interacts with the cytochrome b-c1 complex, cytochrome oxidase complex, heme, Apaf-1, and Caspase 9 proteins. The 6H2.B4 monoclonal antibody recognizes human, mouse, and rat cytochrome-c and has been shown to be useful for intracellular flow cytometric staining, Western blotting, immunoprecipitation, and immunofluorescence staining.

Antigen References: 1. Liu X, *et al.* 1996. *Cell.* 86:147.
 2. Li P, *et al.* 1997. *Cell.* 91:479.
 3. Zhang Z, *et al.* 2003. *Gene* 312:61.
 4. Ferguson H, *et al.* 2003. *J. Biol. Chem.* 278:4579