

Brilliant Violet 605™ anti-T-bet

Catalog # / Size: 3824085 / 125 µl

Clone: 4B10


Isotype: Mouse IgG1, κ

Reactivity: Human, Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 605™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were surface stained with CD3 APC and then treated with True-Nuclear™ Transcription Factor Buffer Set (Cat# 424401). Cells were then stained with T-bet (clone 4B10) Brilliant Violet 605™ (top) or mouse IgG1,


Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining. For flow cytometric staining, the suggested use of this reagent is ≤5 µL per million cells or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use



the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation² and immunofluorescence microscopy³.

NOTE: For flow cytometric staining with this clone, True-Nuclear™ Transcription Factor Buffer Set (Cat. No. [424401](#)) offers improved staining and is highly recommended over the Foxp3 Fix/Perm Buffer Set (Cat. No. 421403) and the Nuclear Factor Fixation and Permeabilization Buffer Set (Cat. No. 422601).

Application References:

1. Szabo SJ, *et al.* 2000. *Cell* 100:655. (ICFC, WB)
2. Hwang ES, *et al.* 2005. *J. Exp. Med.* 202:1289. (ICFC, WB, IP)
3. Neurath MF, *et al.* 2002. *J. Exp. Med.* 195:1129. (IF)
4. Hsieh CY, *et al.* 2012. *J Pharmacol Exp.* 343:125. [PubMed](#).

Description: T-bet, also known as T-box transcription factor T-bet, is considered to be a "master regulator" of Th1 lymphoid development controlling the production of the cytokine IFN- γ . T-bet is widely expressed in hematopoietic cells including stem cells, NK cells, B cells, and T cells. T-bet is critical for the control of microbial pathogens, and knockout animals show multiple physiologic and inflammatory features characteristic of asthma. T-bet expression is optimally observed after IL-12 stimulation and can be suppressed by addition of the Th2 cytokine IL-4 or neutralization of IL-12.

Antigen References:

1. Szabo SJ, *et al.* 2000. *Cell* 100:655.
2. Szabo SJ, *et al.* 2002. *Science* 295:338.
3. Finotto S, *et al.* 2002. *Science* 295:336.
4. Mullen AC, *et al.* 2001. *Science* 292