

**PE/Dazzle™ 594 anti-T-bet**

**Catalog # / Size:** 3824140 / 100 µg  
3824135 / 25 µg

**Clone:** 4B10


**Isotype:** Mouse IgG1, κ

**Reactivity:** Human, Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



Human peripheral blood lymphocytes were surface stained with CD3 APC and then treated with True-Nuclear™ Transcription Factor Buffer Set. Cells were then stained with T-bet (clone 4B10) PE/Dazzle™ 594 (top) or mouse IgG1, κ PE/Dazzle&tra

**Applications:**


**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by intracellular immunofluorescent staining . For flow cytometric staining, the suggested use of this reagent is 1.0 microg per million cells in a staining volume of 100 microL. It is recommended that the reagent be titrated for optimal performance for each application.

\* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>2</sup> and immunofluorescence microscopy<sup>3</sup>.

**NOTE:** For flow cytometric staining with this clone, True-Nuclear™ Transcription Factor Buffer Set (Cat. No. [424401](#)) offers improved staining and is highly recommended over the Foxp3 Fix/Perm Buffer Set (Cat. No. 421403) and the Nuclear Factor Fixation and Permeabilization Buffer Set (Cat. No. 422601).



- Application** 1. Szabo SJ, *et al.* 2000. *Cell* 100:655. (ICFC, WB)
- References:** 2. Hwang ES, *et al.* 2005. *J. Exp. Med.* 202:1289. (ICFC, WB, IP)
3. Neurath MF, *et al.* 2002. *J. Exp. Med.* 195:1129. (IF)
4. Hsieh CY, *et al.* 2012. *J Pharmacol Exp.* 343:125. [PubMed](#).
- 

**Description:** T-bet, also known as T-box transcription factor T-bet, is considered to be a "master regulator" of Th1 lymphoid development controlling the production of the cytokine IFN- $\gamma$ . T-bet is widely expressed in hematopoietic cells including stem cells, NK cells, B cells, and T cells. T-bet is critical for the control of microbial pathogens, and knockout animals show multiple physiologic and inflammatory features characteristic of asthma. T-bet expression is optimally observed after IL-12 stimulation and can be suppressed by addition of the Th2 cytokine IL-4 or neutralization of IL-12.

- Antigen**
- References:** 1. Szabo SJ, *et al.* 2000. *Cell* 100:655.
2. Szabo SJ, *et al.* 2002. *Science* 295:338.
3. Finotto S, *et al.* 2002. *Science* 295:336.
4. Mullen AC, *et al.* 2001. *Science* 292