## **Product Data Sheet**

## PE/Cy7 anti-HA.11 Epitope Tag

**Catalog # / Size:**  $5107635 / 25 \mu g$ 

5107640 / 100 µg

**Clone:** 16B12

**Isotype:** Mouse IgG1, κ

Immunogen: Monoclonal antibody HA.11 was raised

against the twelve amino acid peptide

CYPYDVPDYASL.

Reactivity: Other

**Preparation:** The antibody was purified by affinity


chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7

and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2 mg/ml



CHO-K1 cells (open histogram) or HA tag stably transfected cells (filled histogram) were fixed with Fixation Buffer (Cat. No. 420801), permeabilized with True-Phos™ Perm Buffer (Cat. No. 425401), then intracellularly stained with HA.11 Epitope Tag (

## **Applications:**

**Applications:** Intracellular Flow Cytometry

Recommended

nended Each lot of this antibody is quality control tested by intracellular Usage: immunofluorescent staining with flow cytometric analysis. For flow

immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq\!0.25~\mu g$  per million cells in 100  $\mu l$  volume. It is recommended that the reagent be titrated for optimal performance

for each application.

Application Notes:

Additional tested and reported applications of the 16B12 clone for the relevant

formats include: western blot (WB), immunocytochemistry (ICC),

immunoprecipitation (IP), and flow cytometry (FC).

\*Our Posi-Tag Control Protein (931301) can be used as a helpful positive control

for this antibody.

This second-generation HA antibody is an excellent substitute for the 12CA5 monoclonal antibody. The HA.11 antibody recognizes the influenza hemagglutinin epitope (YPYDVPDYA) which has been used extensively as a general epitope tag

in expression vectors. The extreme specificity of the antibody allows

unambiguous identification and quantitative analysis of the tagged protein. The HA.11 antibody recognizes HA epitopes located in the middle of protein

sequences as well as at the N- or C-terminus.

**Description:** The HA tag (hemagglutinin) is an amino acid seguence derived from the human

influenza hemagglutinin surface glycoprotein, corresponding to amino acids 98-106. It is commonly used as a tag to facilitate detection, isolation, and

purification of proteins. The full amino acid sequence is: YPYDVPDYA.